


# HEADLICE MANAGEMENT CHART



## Treatment Option One: WET COMBING

- Wash the hair in the normal way with an ordinary shampoo
- While the hair is very wet, comb through the hair from the roots with a fine-toothed comb
- Make sure the teeth of the comb slot into the hair at the roots with every stroke.  
This should be done over a pale surface, such as a paper towel or the bath
- Repeat this routine every day for two weeks, so that any lice emerging from the eggs are removed they can be spread

## Treatment Option Two: INSECTICIDES

- Check with your pharmacist, GP or School Nurse which preparations are recommended
- Follow the manufacturer's instructions carefully
- After treatment and whilst the hair is still wet, a fine-toothed comb may be used to remove dead lice
- It could take up to 24 hours for lice to die, so do not assume the treatment has not worked
- To ensure treatment has been successful, detection combing on wet hair should be carried out on all treated persons three times over the next seven days
- REPEAT treatment on the SEVENTH DAY whether or not lice are found
- Small lice means that the eggs were not affected by the first treatment. A repeat treatment will kill the baby lice before they can lay more eggs
- Large lice means that the head has been re-infested – so contact tracing and the treatment cycle will have to be restarted

## AFTER SUCCESSFUL TREATMENT:

↓  
Comb wet hair using a fine-toothed comb at least Once a week

## IF TREATMENT NOT SUCCESSFUL

↓  
IF LARGE LICE FOUND:  
Check family and friends for Headlice

↘  
IF SMALL LICE FOUND:  
Use Treatment One

**ACTION:**  
Repeat treatments

If treatment Option One used previously use Treatment Option Two

If treatment Option Two used previously use Treatment Option One